

**Michigan**  
**Adult Cardiac Protocols**  
**ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA)**

Date: May 31, 2012

Page 1 of 1

## ***Asystole / Pulseless Electrical Activity***

During CPR, consider reversible causes of Asystole/PEA and treat as indicated. Causes and efforts to correct them include but are not limited to:

Hypovolemia – give NS IV/IO fluid bolus up to 1 liter, wide open.

Hypoxia – reassess airway and ventilate with high flow oxygen

Tension pneumothorax – see **Pleural Decompression Procedure**

Hypothermia – follow **Hypothermia Cardiac Arrest Protocol** rapid transport

Hyperkalemia (history of renal failure) – see #5 below.

### **Pre-Medical Control**

#### **PARAMEDIC**

1. Follow the **Cardiac Arrest - General Protocol**.
2. Confirm that patient is in asystole by evaluating more than one lead.
3. Administer Epinephrine 1:10,000, 1 mg (10 ml) IV/IO, repeat every 3-5 minutes.
4. Per MCA selection, administer Vasopressin 40 Units IV/IO in place of second dose of Epinephrine.

**Vasopressin**

40 Units IV/IO

Included

Not Included

5. In a dialysis patient hyperkalemia is likely. Administer Calcium Chloride 1gm IV/IO and Sodium Bicarbonate 1 mEq/kg IV/IO with 20 ml NS flush in between medications.
6. Continue CPR and reassess rhythm every 2 minutes.