

Genesee County
General Procedures
VASCULAR ACCESS & IV FLUID THERAPY

Date: February 2, 2017

Page 1 of 4

Vascular Access & IV Fluid Therapy

Purpose: To outline the process in patients requiring vascular access. This policy applies to Specialists and Paramedics.

Indications

1. For the purpose of fluid or medication administration.
2. External jugular cannulation should be initiated in patients in whom access is necessary and other peripheral vascular access is not accessible or is contraindicated.
3. IO indications: Adult and pediatric life threatening situations where venous access using peripheral veins has been unsuccessful. IO access should be considered early in situations where IV access is unsuccessful or technically challenging. Indications include:
 - A. Cardiac Arrest
 - B. Severe burn injury with shock
 - C. Shock
 - D. Severe multiple trauma with shock
 - E. Status epilepticus
 - F. Contact medical control for other situations without delaying transport

Saline Lock may be initiated in patients in whom IV access for medication administration may be necessary but IV fluid therapy unlikely.

IVs will be initiated in those situations in which fluid resuscitation may be indicated.

Contraindications

1. To peripheral vascular access:
 - A. No peripheral sites available
 - B. Burns overlying available peripheral sites unless no other sites available
 - C. Infection overlying available peripheral sites
2. To intraosseous infusion and placement:
 - A. If infiltration occurs (rare), do not reuse the same bone as fluid will leak out of the original hole; select another site.
 - B. Do not place in a fractured extremity. If the femur is fractured, use the opposite leg.

Special Considerations (Side effects/Complications)

1. Initiation of vascular access generally should not delay patient transport to the hospital.
2. General side effects or complications: infection, air embolism, catheter shear, hematoma, arterial puncture, fluid overload
3. Intraosseous placement:

Genesee County
General Procedures
VASCULAR ACCESS & IV FLUID THERAPY

Date: February 2, 2017

Page 2 of 4

- A. Complications include subperiosteal infusion, osteomyelitis, sepsis, fat embolism, bone marrow damage.

Standards for IV attempts

1. Two (2) attempts per provider, maximum 4 attempts.
2. Consider IO early, as indicated above.
3. Document any reasons for deviation.
4. If the patient has a PICC line, the paramedic may use the PICC line for vascular access on any Priority 1 patient or with online medical control approval and bypass the other access attempts noted above. A PICC line can only be used for fluids and medications.
 - A. All cases where a PICC line has been accessed or was attempted to be accessed must be submitted to the GCMCA for review by the 15th of the following month.
 - B. All agencies will provide training to all paramedics on accessing PICC lines consistent with an outline created by the GCMCA Education Committee.

Needle size for IV placement

1. Adult TKO 18 ga - 20 ga Angiocath
2. Adult trauma, bleeding or cardiac arrest 14 ga - 18 ga.
3. Pediatrics 20 ga - 24 ga Angiocath

Flow Rates

1. Flow rates for all IV's are to be at rates TKO or saline lock unless otherwise indicated by specific protocol or Medical Control.
2. The amount of fluid infused along with the IV rate is to be noted on the EMS Medical Record
 - A. 25 ml/hr is TKO rate.
 - B. Saline lock IV is preferred in place of TKO IV lines.
3. Flow rates and changes in flow rates must be documented on the EMS Patient Care Record.
4. The standard IV/IO fluid bolus volume will be 1 liter normal saline with repeat as necessary, unless otherwise noted by protocol. IV/IO fluid bolus is contraindicated in patients with pulmonary edema. Volume for pediatric IV/IO fluid bolus is 20 ml/kg, unless otherwise noted by protocol.
5. Medicated drips should be piggybacked into the main IV line or saline lock.

Solutions – Unless otherwise specified the IV solution of choice is Normal Saline 0.9% (NS).

IV Tubing

1. Normal Saline – macrodrip
2. Children - macrodrip

Genesee County
General Procedures
VASCULAR ACCESS & IV FLUID THERAPY

Date: February 2, 2017

Page 3 of 4

Procedure IV/IO Placement

1. Utilize universal precautions for all IV/IO placements.

Procedure for Peripheral Vascular Cannulation:

1. Gather and prepare equipment.
2. Place the tourniquet on the extremity.
3. Cleanse the skin
4. Make your puncture while maintaining vein stability.

5. Watch for flashback. Once you have a blood return, advance the catheter as per normal IV technique and attach the IV tubing or Normal saline lock tubing and cap.
6. If you have no blood return and you are in the vein, remove the needle hub and attach your syringe to assist in aspirating for blood.
7. Instill 2-3 ml of normal saline if normal saline lock placed.
8. Secure catheter and IV tubing.

Procedure for External Jugular Cannulation:

1. Gather and prepare equipment
2. Position patient supine (trendelenburg, if possible)
3. Turn head to opposite side of venipuncture (if no C-spine injury is suspected)
4. Cleanse the skin
5. Occlude the vein by using the side of your finger above the clavicle to facilitate filling the vein.
6. Make your puncture midway between the angle of the jaw and the middle of the clavicle.
7. Watch for flashback. Once you have a blood return, advance the catheter as per normal IV technique and attach the IV solution or normal saline lock cap, covering catheter with gloved finger while preparing to attach the IV tubing. If you have no blood return and you are in the vein, remove the needle hub and attach your syringe to assist in aspirating for blood.
8. Instill 2-3 ml of normal saline if normal saline lock placed.
9. Secure IV catheter and tubing.

Procedure for Intraosseous Placement:

1. Have all IO equipment ready prior to bone penetration.
2. Expose the extremity.
3. Stabilize the extremity to minimize motion.
4. Selection of site:
 - A. Proximal Tibia or Proximal Humerus.
 - B. In children less than six years of age, the preferred site is the proximal tibia.
5. Insertion:
 - A. Follow the manufacturer's recommendations for IO insertion with the

Genesee County
General Procedures
VASCULAR ACCESS & IV FLUID THERAPY

Date: February 2, 2017

Page 4 of 4

-
- above indications.
6. Scrub the insertion site with alcohol prep/chlorhexidine. Strict adherence to aseptic technique is essential.
 7. Insert the IO needle.
 8. Attempt to confirm marrow placement by removing the stylet and aspirating blood and/or bone marrow.
 - A. If unable to aspirate, attach 10 – 20 ml syringe with normal saline and gently infuse normal saline.
 - B. Observe for normal saline leakage or SQ tissue swelling.
 - a. If neither occurs, proceed.
 - b. If either occurs, select a different site.
 9. Connect the appropriate IV equipment (normal saline locks not indicated in IO placement).
 10. Administer the appropriate fluids and/or drugs.
 11. Stabilize the entire intraosseous set-up as if securing an impaled object.
 12. In conscious patients experiencing pain with IO infusion consider administering Lidocaine 2 %, 20 mg IO for adult patients, 0.5 mg/kg for pediatrics administer to a maximum of 20 mg. (Lidocaine 2% = 20 mg/ml).
 13. Notify Medical Control of the IO placement.
 14. If the IO is unsuccessful after 2 attempts, contact Medical Control.