

Michigan MEDICATION SECTION DIPHENHYDRAMINE

Initial Date: 07/19/2023 Revised Date:

Diphenhydramine

Pharmacological Category: Histamine H1 Antagonist

Routes: IV/IO/IM

Indications:

- 1. Anaphylaxis
- 2. Mild or moderate allergic reaction
- 3. Urticaria/hives
- 4. Nausea and vomiting

Expected effects:

- 1. Antihistamine, decreased urticarial, decreased itching
- 2. Drowsiness

Dosing: NAUSEA AND VOMITING

Indications: Nausea and vomiting <u>Adults</u> administer:

1. Diphenhydramine 12.5-25 mg IV/IM. Maximum dose 25 mg.

<u>Pediatric</u> (>2 years of age AND > 12 kg) administer:

- 1. According to MI MEDIC cards
- 2. If MI MEDIC cards are not available administer:
 - a. Diphenhydramine 1.0 mg/kg IV. Max dose 25 mg.

Dosing: ANAPHYLAXIS ALLERGIC REACTION

Indication: Anaphylaxis/allergic reaction <u>Adults</u> administer:

1. Diphenhydramine 50 mg IM/IV/IO

Pediatrics administer:

- 1. According to MI MEDIC cards
- 2. If MI MEDIC cards are not available administer:
 - a. Diphenhydramine 1 mg/kg IM/IV/IO. Maximum dose 50 mg.

Dosing: POISONING/OVERDOSE/ENVIRONMENTAL EXPOSURE

Indication: extrapyramidal dystonic reactions <u>Adults</u> administer:

1. Diphenhydramine 50 mg IV.

Pediatrics administer:

1. Diphenhydramine 1 mg/kg IV. Max dose 50 mg.



Michigan MEDICATION SECTION DIPHENHYDRAMINE

Initial Date: 07/19/2023 Revised Date:

<u>Used in the Following Protocols</u> Nausea & Vomiting (Section 1 General Treatment) Anaphylaxis/Allergic Reaction (Section 1 General Treatment) Poisoning/Overdose/Environmental Exposure (Section 2 Trauma and Environmental)