Genesee County Medical Control Authority Medication Protocols PHARMACY

February, 2024 Section 9-6

Pharmacy

Medication Exchange and Replacement Procedure

VEHICLE STOCK

- A. Each approved ALS unit will carry one GREEN LOCK SEALED Southeast Michigan (SEM) Regional Medication Box and A-Pack (Ancillary Pack). Contents are listed in Pharmacy Appendixes 1 and 2. Only appropriately numbered Medication Boxes and A-Packs issued by the participating Medical Control Authority are to be stocked by participating hospital pharmacies and issued to approved ALS units.
- B. Each EMS agency will be responsible for providing any additional equipment required by Michigan Department of Health & Human Services Bureau of EMS & Trauma (MDHHS).
- C. All drugs, needles, syringes, and supplies will be stored in a securely locked and temperature controlled location on each approved unit. Medication Boxes/A-Packs will remain sealed at all times except when in actual use.
- D. Medication Boxes/A-Packs are to be inspected daily by the crew of the unit for evidence of loss, theft, discrepancy, and expiration date. Inspection items include, but are not limited to: the Medication Box/A-Pack is locked in a compartment, the green lock is intact, the lock # matches number on the label, and medications are not expired. It is recommended that this inspection be included in a standard documented vehicle checklist.
- E. Agencies are responsible for maintaining Medication Boxes/A-Packs not in use by a crew. At a minimum the boxes must be secured in a locked area or cabinet that can only be accessed by paramedics or agency leadership. The area where the boxes are located must have video surveillance that is capable of capturing any activity with the boxes. Agencies must also have a check-in/check-out process that paramedics use to obtain and return the boxes.
- F. Unopened Medication Boxes/A-Packs are to be exchanged within seven (7) days of the, "Use or Replace By" date.

USE/REPLACEMENT/EXCHANGE

- A. Medication Boxes/A-Packs will only be opened by a Paramedic when presented with a patient requiring Advanced Life Support care (when acting on written or transmitted orders from a physician at an appropriate On-Line Medical Control Facility) or the Pre-Medical Control section of approved treatment protocols.
- B. Red/Green Lock Procedure for Medication Boxes/A-Packs
 - 1. The Medication Box/A-Pack will be sealed using a green lock bearing the number indicated on the label.
 - 2. After the pharmacy inventory/restocking is complete, a red lock bearing the number indicated on the label will be placed in the Medication Box/A-Pack to be used by the Paramedic to seal the Medication Box/A-Pack after it has been used.
 - 3. When the Medication Box/A-Pack is opened by the Paramedic the broken numbered green lock will be placed in the Medication Box/A-Pack and delivered with the used Medication Box/A-Pack to the replacing pharmacy.

Medication Protocols PHARMACY

February, 2024 Section 9-6

- 4. After use the Paramedic will seal the Medication Box/A-Pack for exchange with the red lock from the Medication Box/A-Pack bearing the number indicated on the label.
- C. <u>OPTIONAL (MCA adoption required)</u> Red/Green/White/ (or Yellow) Lock Procedure for MEDICATION BOXES ONLY
 - 1. After the pharmacy inventory/restocking is complete, a red lock and green lock bearing the respective numbers indicated on the label will be placed in the Medication Box to be used to seal the box after initial inspection (green lock) and after post use inspection (red lock).
 - 2. The Medication Box will be sealed using a white (yellow) lock.
 - 3. After the Medication Box is inspected jointly by the Paramedic and ED/Pharmacy representative the Medication Box will be sealed with the green lock, from the Medication Box, bearing the number indicated on the label.
 - 4. When the Medication Box is opened by the Paramedic, the broken numbered green lock will be placed in the Medication Box and delivered with the used Medication Box to the replacing pharmacy. If only an A-Pack is used in treating a patient and that patient refuses transportation, the crew should go to the nearest hospital to exchange the A-Pack but will remain in service should they be needed for another emergency call. If they are dispatched to another emergency call prior to exchanging the A-Pack they can delay the exchange to respond to that call and exchange the A-Pack at their next opportunity.
 - 5. After use, and after joint inspection of the Medication Box for exchange by the Paramedic and ED/Pharmacy representative, the Paramedic will seal the Medication Box with the red lock from the Medication Box bearing the number indicated on the label.

MEDICATION BOXES:

- A. All Participating Hospitals will have Medication Boxes/A-Packs, with contents as approved by the participating Medical Control Authorities and MDHHS, available for replacement of supplies used by approved ALS Units. Replacement Medication Boxes/ A-Packs will be maintained in a locked area, under the control of hospital staff, which is available 24 hours a day, 7 days a week. This area will be located within the either Emergency Department or Pharmacy of the Participating Hospital. Appropriate record keeping and security measures are required at each exchange site to ensure that only appropriately licensed and authorized personnel have access to medications and other related supplies.
- B. Medication Boxes/A-Packs used by approved ALS units for patients transported will be replaced, at the time of the run, by the receiving hospital according to established procedure. Where the receiving facility does not participate in the Regional EMS Medication Exchange System and/or supplies are expended for a patient who subsequently is not transported, the unit will proceed immediately to the Regional Participating Hospital which provided Medical Control for the run to complete replacement. A PCR will be submitted when completed.
- C. Use of any supplies contained in the Regional Medication Box/A-Pack will be

Medication Protocols PHARMACY

February, 2024 Section 9-6

documented on the Use/Replacement Form for exchange and the PCR of the patient for whom the supplies were used. This includes any medications or supplies prepared for use but not actually administered to the patient.

Medication Protocols PHARMACY

February, 2024 Section 9-6

BOX CLEANING

- A. All empty containers, packaging and used materials will be properly disposed of by the ALS crew that used the Medication Box/A-Pack.
- B. The EMS crew using standard hard surface decontamination techniques will clean any blood or body fluid contamination to the exterior of the Medication Box.
- C. If there is blood or body fluid contamination to the interior of the Medication Box/A-Pack, or to any unused materials or packaging, the EMS crew will clean and dispose of contaminated material. If direction is needed in the cleaning and disposal of contaminated materials the crew can contact the receiving hospital pharmacy.
- D. All unused, un-contaminated supplies will be returned to the Medication Box/A-Pack.

THE ALS CREW WILL:

- A. For all SEM runs, complete the Use/Replacement Form contained in the Medication Box/A-Pack. The form shall serve as the permanent medical record for drugs administered, and the paramedic will document their MCA Medical Director's name when any controlled substance is used. For post-radio controlled substance orders, document the ordering physician's name on the Replacement Form.
- B. The ALS crew is responsible for proper distribution of the completed forms.
- C. The expended Medication Box/A-Pack (cleaned as described above and red sealed) and the completed Use/Replacement Form will be presented to an appropriate member of the hospital staff who will issue a fresh Medication Box/A-Pack (green seal). A member of the ALS crew and the hospital staff member will complete the exchange log sheet.
- D. In the event that controlled substances are prepared for use and not used or the entire contents of a container are not used, the remaining medication will be appropriately wasted by ALS crew member in the presence of licensed hospital personnel/or other ALS crew member. The following will be recorded on the Use/Replacement Form:
 - 1) The name and amount of the medication wasted.
 - The initials of the ALS crew member, hospital personnel or other ALS crew
 - member witnessing the waste.
- E. All requests for information concerning the "Use/Replacement Form" by other agencies are to be directed to the appropriate Medical Control Authority.

EXPIRATION OF DRUGS/SOLUTIONS

- A. All items in a SEM Regional Medication Box/A-Pack will have expiration dates not less than 90 days after the Medication Box/A-Pack is prepared.
- B. Any unused items bearing expiration dates less than 90 days subsequent shall be removed from the Medication Box/A-Pack and replaced with fresh stock as described in A above.
- C. Each Regional Medication Box/A-Pack will have a label securely attached to the

Medication Protocols PHARMACY

February, 2024 Section 9-6

outside containing the following information:

- 1. The name of the participating hospital pharmacy, which restocked the Medication Box/A-Pack.
- The date the Medication Box/A-Pack was restocked.
- 3. The printed name and initial of the pharmacist and pharmacy technician that inventoried and restocked the Medication Box/A-Pack.
- 4. The expiration date is the last day of the month of the earliest expiring medication (with a maximum of one year from the current date). The Medication Box/A-Pack label will include the month/day/year in the "use or replace by" section.
- 5. The red and green lock numbers.
- 6. The Medication Box/A-Pack number.

MEDICATION BOXES - ALTERNATIVE PACKAGING AND SHORTAGES:

- A. Routinely, participating hospital pharmacies must provide items only in the dosage, concentration, and packaging listed. Use of alternative vendors or manufacturers is acceptable if consistent with the required contents.
- B. For products in short supply hospital pharmacies may stock the Medication Boxes/A- Packs with less than a 90-day expiration date.
- C. When a medication in alternative packaging is the only product available, place alternative medication, use directions and supplies for medication preparation inside the
 - Medication Box/A-Pack.
- D. Attach a sticker to the exterior top of the Medication Box or to the clear side near the bottom of the A-Pack stating the substitution.
- E. Directions for specific medications in short supply, throughout the regional exchange system will be addressed through communications with participating pharmacies as approved by the Regional Protocol participating MCAs.

DISCREPANCIES

- DEFINITION: For purposes of this policy, a "discrepancy" is any breakage, expiration, shortage, theft or diversion of a Regional Medication Box/A-Pack, or any contents thereof.
- A. A standard "MEDICATION DISCREPANCY REPORT" will be completed each time a discrepancy occurs. The form may be initiated by either pre-hospital or hospital staff discovering the discrepancy. The person initiating the report will be responsible for distributing the forms as required.
- B. The Medical Control copy of discrepancy reports will be sent to the Medical Control Authority in which the discrepancy occurred, which will serve as the central filing point.
- C. A copy of the PCR for the run on which the discrepancy occurred/was discovered is to be attached to each copy of the discrepancy report where applicable.
- D. The participating hospital pharmacist is to be notified immediately if controlled substances are involved in a discrepancy. The participating hospital pharmacist

Medication Protocols PHARMACY

February, 2024 Section 9-6

will determine if the discrepancy constitutes a diversion of controlled substances. In addition, the following are to be notified of controlled substance diversions:

- 1. The Medical Control Authority in which the diversion occurred.
- 2. Drug Enforcement Administration (DEA)
- 3. Michigan State Board of Pharmacy
- 4. Appropriate local law enforcement agency (for the jurisdiction where the diversion most likely took place)
- 5. Michigan Department of Health & Human Services (MDHHS).
- E. <u>The participating hospital pharmacist will be responsible for assuring that</u> all appropriate notifications are made.
- F. If, at any time, an ALS unit has less than the required stock of Medication Box/A-Pack supplies and cannot document use of these supplies in connection with a patient, a discrepancy report must be completed. The completed discrepancy report, along with a completed Use/Replacement Form indicating the EMS Provider Agency Name under "Patient Name" and clearly marked "Replacement for Missing Stock" will be presented to the agency's Base Hospital Pharmacy for replacement. The ALS agency can be held accountable for replacement.

Medication Protocols PHARMACY

February, 2024 Section 9-6

| rebruary, 2024 | | Section 9-6 | | | |
|--|---------------------------|---------------------------------------|-------------------|--|--|
| MEDICATION BOX CONTENTS DRUG/ITEM | CONCENTRATION | PACKAGING | QUANTITY | | |
| Acetaminophen | 650 mg/20.3 mL | Unit Dose Cup | 1 | | |
| Adenosine | 6 mg/2 mL | 2 mL Vial/Syringe | 3 | | |
| Albuterol | 2.5 mg/3 mL | 3 mL Vial - UD | 6 | | |
| Amiodarone | 150 mg/3 mL | Amp/Vial | 3 | | |
| Aspirin | 81 mg/tablet | BT/UD – chewable | 1 BT or 4 UD tabs | | |
| Atropine | 1 mg/10 mL | 10 mL Syringe | 3 | | |
| Calcium Chloride | 1 g/10 mL | 10 mL Syringe | 2 | | |
| Ceftriaxone | 2gm vial | 2gm vial | 1 | | |
| Dextrose 50% | 25 g/50 mL | 50 mL Syringe | 1 | | |
| Diphenhydramine | 50 mg/1 mL | 1 mL Vial | 2 | | |
| Epinephrine | 1 mg/1 mL | 1 mL Amp/ Vial | 2 | | |
| Epinephrine | 1 mg/10 mL | 10 mL Syringe | 7 | | |
| Fentanyl | 50 mcg/mL | 2 mL Vial/Amp | 3 | | |
| Ipratropium Bromide | 0.02% | 2.5 mL Vial - UD | 2 | | |
| Ketamine | 100mg/ml | 5ml Vial | 1 | | |
| Ketorolac | 15mg/ml | 1ml Vial | 1 | | |
| Lidocaine | 100 mg/5 mL | 5 mL Syringe | 3 | | |
| Magnesium Sulfate | 1 g/2 mL | Amp/Vial | 4 | | |
| Methylprednisolone | 125 mg | Vial | 1 | | |
| Midazolam | 5 mg/1 mL | 1 mL Vial | 4 | | |
| Morphine | 10 mg/1 mL | 1 mL Amp/Vial | 2 | | |
| Naloxone | 2 mg/2 mL or 0.4 mg/mL | 4 x 2 mL Syringe or 2 x 10 mL Vial | Total = 8mg | | |
| Nitroglycerin | 0.4 mg/tab | Bottle | 1 | | |
| Ondansetron | 2 mg/mL | 2 mL Vial | 2 | | |
| Ondansetron ODT | 4mg | Tablet | 2 | | |
| Prednisone | 50 mg tab | 50 mg Tab | 1 | | |
| Racepinephrine 2.25% with 3 mL NS | 11.25 mg/0.5 mL | 0.5 mL Vial | 1 | | |
| Sodium Bicarbonate | 50 mEq/50 mL | 50 mL Syringe | 2 | | |
| Sodium Chloride | 0.9% | 100 mL Bag | 1 | | |
| Sodium Chloride | 0.9% Preservative Free | 20-30 mL Vial or 10 mL syringes | 1 2 | | |
| Tranexamic Acid (TXA) | 100mg/ml | 10 ml Vial | 1 | | |
| Alcohol Pad | | | 12 | | |
| Incident Report Form | | | 1 | | |
| IV Additive Labels | | | 3 | | |
| IV Tubing with Y Site Pre-pierced Reseal | 60 drops/mL(mini drip) | | 2 | | |
| Nebulizer | | | 1 | | |
| Blunt Cannula | 18 G x 1 inch | | 5 | | |
| Filter Needle | 18-21 G | | 3 | | |
| Intranasal Mucosal Atomization Device | | | 1 | | |
| Syringe | 20 mL | | 1 | | |
| Syringe | 10 mL | | 5 | | |
| Syringe with needle/Luer Lock | 1 mL | | 5 | | |
| Syringe with needle | 3 mL - 21/22 G x 1.5 inch | | 5 | | |
| Oral Liquid Syringe | 10 ml | | 1 | | |
| Needle | 18 G x 1.5 inch | | 3 | | |
| Pediatric Needle | 25 G x 1 inch | | 2 | | |
| Red Lock | | | 1 | | |
| Replacement Form | | | 1 | | |
| Three or Four-Way Stopcock | | | 1 | | |
| · · · | | | • | | |

NOTE: Participating hospital pharmacies must provide the above listed items only in the dosage, concentration, and packaging shown above. Use of alternative vendors or manufacturers is acceptable if consistent with the required contents.

SEM/EMS MEDICATION BOX CONTENTS AND SCHEMATIC

| To | n | Sh | el | lf |
|-----|---|----|----|----|
| - U | • | | | |

| Acetamin 650 mg/ 20 | | Sodium Chl Preservative Free | | Misc. Su Alcohol Pa | | |
|---------------------------------------|--------------|--|---|--|--|--|
| Unit dose of | | Vial or (2) 10 mL | | Blunt Cannula (18 G x 1 inch) – x 5 | | |
| | | Nalo 2mg/ 2ml Syring | <u>xone</u> | Filter Needle 18 IV Additive Needle (18 G x Pediatric Needle (2 Three or Four Wa | 8 - 21 G - x 3 Labels x 3 1.5 inch) - x 3 25 G x 1 inch) x 2 ay Stopcock x 1 | |
| Magnesium 1 g/2 1 Amp/ V X 4 | mL ⁄ial | 2 mg/ 2 mL o 4 x 2 mL Syringe Total = Intranasal Mucosal At | r 0.4 mg/ mL or 2 x 10 mL Vial = 8 mg | Red Lo | ck x l | |
| <u>Amiodarone</u> | Adenosine | Epinephrine | Diphenhydramine | Aspirin 81 mg | Ondansetron | |
| 150 mg/ 3 mL | 6 mg/2 mL | 1mg/ 1 mL | 50 mg/ 1 mL | Chewable | 2 mg/ mL | |
| Amp/ Vial | 2 mL | Amp/ Vial | 1 mL Vial | <u>Tablet</u> | 2 mL Vial | |
| X 3 | Vial/Syringe | X 2 | X 2 | X 1 Bottle OR 4 | X 2 | |
| | X 3 | | | UD Tabs | | |
| | | | Tranexamic Acid | Nitroglycerin | Ondansetron | |
| | | | 100mg/ml | 0.4 mg/ Tab | 4mg ODT | |
| | | | 1 x 10ml vial | Bottle | 2 Tabs | |
| Middle Chale | | | | X 1 | | |

Middle Shelf

| Controlled Substances | Methylprednisolone | <u>Ipratropium</u> | Albuterol | <u>Nebulizer</u> |
|-----------------------|--------------------|--------------------|----------------|-----------------------|
| <u>Fentanyl</u> | 125 mg/ Vial | Bromide 0.02 % | 2.5 mg/ 3 mL | X 1 |
| 50 mcg/ mL - | X 1 | 2.5 mL Vial – UD | 3 mL Vial – UD | |
| 2 mL Vial/Amp x 3 | | X 2 | X 6 | Racepinephrine |
| <u>Midazolam</u> | Prednisone | | | 2.25 % |
| 5 mg/ 1 mL- | 50 mg Tablet | Ketorolac | | 11.25 mg/ 0.5 mL |
| 1 mL Vial x 4 | X 1 | 15mg/ml Vial | | 0.5 mL Vial |
| <u>Morphine</u> | | X 1 | | X 1 |
| 10 mg/ 1 mL- | <u>Ceftriaxone</u> | | | 3 mL NS |
| 1 mL Vial/Amp x 2 | 2gm vial | | | X 1 |
| <u>Ketamine</u> | X 1 | | | |
| 100mg/ml | | | | |
| 5ml Vial x 1 | | | | |

Bottom Shelf

| Bag of Syringes | Sodium Bicarbonate | Epinephrine |
|--|---------------------------------|-----------------------------------|
| Syringe (With needle/ Luer Lock) – 1 mL x 5 | 50 mEq/ 50 mL – 50 mL | 1 mg/ 10 mL – 10 mL Syringe x 7 |
| Syringe 3 mL (21/22 G x 1.5 inch) – 3 mL x 5 | Syringe x 2 | |
| Syringe – 10 mL x 5 | | |
| Syringe – 20 mL x 1 | Dextrose 50% | |
| | 25 g/ 50 mL – 50 mL Syringe x 1 | IV Tubing With Y Site Pre-pierced |
| Lidocaine | | Reseal |
| 100 mg/5 mL - 5 mL Syringe x 3 | | - 60 drops/mL (mini drip) x 2 |
| | | |
| Calcium Chloride | | |
| 1 g/ 10 mL – 10 mL Syringe x 2 | | |
| | | <u>Forms</u> |
| <u>Atropine</u> | | Replacement/ Schematic/ Incident- |
| 1 mg/ 10 mL – 10 mL Syringe x 3 | | Discrepancy |
| | | |
| Sodium Chloride | | |
| 0.9 % - 100 mL Bag x 1 | | |
| | | |

SEM/EMS ACCESSORY PACK (A-PACK) CONTENTS

Version: February 1, 2024 (Discard all previous versions) Needleless stock only!

| DRUG/ITEM | CONCENTRATION | PACKAGING | QUANTITY |
|---------------------------------------|---------------------------|--------------------------------|--------------|
| Albuterol | 2.5 mg/ 3 mL | 3 mL Vial – UD | 6 |
| Aspirin | 81 mg/Chewable tablet | UD Tabs | 4 |
| Dextrose 50% | 25 g/50 mL | 50 mL Syringe | 1 |
| Intranasal Mucosal Atomization Device | | | 1 |
| Ipratropium Bromide (in baggie) | 0.02% | 2.5 mL Vial – UD | 1 |
| Naloxone | 2 mg/2 mL or 0.4 mg/mL | 2x2 mL Syringe or 1x10 mL Vial | Total = 4 mg |
| Nitroglycerin | 0.4 mg/ Tab | Bottle | 1 |
| Nebulizer | | | 1 |
| Ondansetron | 2 mg/ mL | 2 mL Vial | 2 |
| Ondansetron ODT | 4mg | Tablet | 2 |
| Prednisone | 50 mg tab | 50 mg Tab | 1 |
| Blunt Cannula | 18 G – 1 inch | | 2 |
| Syringe 3 mL with needle | 21/22 G x 1.5 inch needle | | 2 |
| Red Lock | | | 1 |
| Replacement Form | | | 1 |
| Incident Report Form | | | 1 |
| Three or Four-Way Stopcock | | | 1 |

SEM/EMS ACCESSORY PACK (A-PACK) SCHEMATIC

Green Lock through zipper and eyelet

| (Place behind Albuter Dextrose 50% 50 mL Syringe 25 gm/ 50 mL (1) | ol on this side) | Nebulizer (Place on th | | (Elastic Holder) Nitroglycerin 0.4 mg/ Tab (1) bottle |
|--|------------------------------------|---|---|--|
| (Lunida Funud Danland | ` | Incident Repor | t Form (1) | (1) bottle |
| (Inside Front Pocket Albuterol 2.5 mg/ 3 mL | Blunt Cannula 18 G x 1 inch (2) | Replacement F | | 1. (A.B. 1) |
| Vial UD (6) | | (Folded in half | and placed along inside bac | ck of A-Pack) |
| | | Ipratropium Bromio | de 0.02% Vial | |
| Prednisone (In baggie) (1) | | xone 2 mg/ 2 mL or 0.4 mg 2 mL Syringe 1 x 10 mL Total = 4 mg | | 50 mg Tab UD (1) |
| Aspirin 81 mg Tab UD Chewable (4) | Y | (Inside Front Pocket) Yellow Pharmacy Label | Three or Four-Way Stope | cock (1) |
| | Syringe | e 3 mL with 21 G x 1.5 inc. | h needle (2) | Red Lock (1) |
| Intranasal Mucosa | al Atomization Device (1) | Ondansetron 2 mg/ ml | L - 2 mL vial (2) Ondanset 2 Tablets | ron ODT 4mg |

SOUTHEAST MICHIGAN (SEM) REGIONAL MEDICATION BOX/A-PACK AND IV EXCHANGE PROCEDURES

PLEASE POST IN ALL MEDICATION EXCHANGE AREAS

- STEP 1: EMS Personnel must complete a SEM Med Box/A-Pack/IV Supply Use/Replacement Form and/or the SEM IV Supply Use/Replacement Form (EMS Run Report Genesee County MCA). All information must be complete. Used Medication Boxes/A-Packs must be cleared of contaminated items, cleaned, and sealed appropriately.
- STEP 2: Hospital staff reviews form for completeness and receiving prescriber signature (only required for cases in which controlled substances are used). Staff unlocks cabinet and allows removal of appropriate supplies. Both EMS personnel and hospital staff complete the Medication Box/A-Pack and IV Supply Exchange Log. Both EMS and hospital staff ensure that the correct Medication Box/A-Pack numbers are recorded.
- STEP 3: The original copy of the SEM Medication Box/A-Pack/IV Supply Use/Replacement Form shall be left in the MCA cabinet. Because the hospital staff person must review the documentation form, it may not be able to be placed in the Medication Box/A-Pack before it is sealed. It will be necessary for the pharmacist to collect all separated Documentation Logs that are stored in the cabinet, when restocking drug boxes.
- **STEP 4:** The MCA cabinet must be re-locked when the exchange is complete.

THESE PROCEDURES ALSO APPLY WHEN ONLY AN IV FLUID/SUPPLY EXCHANGE IS COMPLETED.

NOTE: Receiving Prescriber: Physician, P.A., N.P.

| Genesee | County | Medica | l Contro | Author | ity P | harn | nacy | Log | | | | 0 | | | nust be con | npleted | | return Green or Red | | | |
|-----------------------------|------------|------------------------|---------------|-------------|-------|-------|---------|--------|--------|--------|--------|-----|------------------------|---------------|-------------|---------------------------|-----------------------|--|----------------|------------|------------------------|
| Hospital _ All columns N | ALIOT DE C | OMDI ETE | 0 | | | | | _ | | | | | for the | following: | | | | issing, broken or tam Blank cells on this R | | | |
| All COlumns I | IUST BE C | OWPLETE | U | | | | | | | | | | | | | | 3. | BIATIK CEIIS ON UNIS R | eport Form | | |
| Date and | Time | | IN | | | Contr | olled : | substa | ance s | status | | | | 0 | UT | | | Personi | nel | | > |
| Date | Time | Box / A-Pack No. | Green Lock | Red Lock | 2 | 3 | conta | 1 | (| On rui | n shee | et | Box / A-Pack No. | Green Lock | Red Lock | All C.S Present Y/N | Pharmacy Personnel | EMS Provider Last Name - Please Print | EMS Badge # | EMS Agency | Discrepency Report? |
| | | | | | Mo | F | Mid | Ket | Мо | F | Mid | Ket | | | | | | | | | |
| 11/1/2013 | 6:45am | 123 | 12345 | 56789 | 2 | 2 | 4 | 1 | | 1 | - | | 456 | 23456 | 67891 | | JT | M. Smith | 1 | ABC | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | |

= Record # of containers present and used. Initiate a Discrepancy Form if sum is not 10. Mo = Morphine 10mg, F = Fentanyl 2 ml, Mid = Midzolam 5 mg, Ket = Ketamine 5 ml

SEM MED BOX/A-PACK SUPPLY USE/REPLACEMENT FORM Version February 1, 2024

| AGENCY/UNIT #: | | | | EMO OF | EW (Names): | | | | |
|--|--------------------------------|--------------|---------------|----------------|--|----------------------------------|------------|------------|-----------|
| NCIDENT #: | | | | | EW (Names): | | | | |
| atient Name: | | | | Patient | OOB: | | | | |
| MEDICATION | UNIT/SIZE | QNTY | USED | NOTE | Needleless stock only! | * Items in both | | | |
| Acetaminophen 650 mg/20.3 mL | Unit dose cup | 1 | | | MISCELLANEOUS Alcohol Pads | UNIT/SIZE | QNTY 12 | USED | NOTE |
| 10 ml oral syringe in bag | | | | | | 4 P 1 | | | |
| Adenosine 6 mg/2 mL | Vial/Syringe 2 mL | 3 | | | Incident Report Form* | A-Pack | 1 Each | | |
| Albuterol 2.5 mg/3 mL* | Vial – UD 3 mL | 6 | | | IV Additive Labels | | 3 | | |
| | A-Pack | 6 | | | IV Tubing 60 drops/mL | | 2 | | |
| Amiodarone 150 mg/3 mL | Amp/Vial | 3 | | | (Minidrip) with Y Site | | | | |
| Aspirin 81 mg chewable | X 1 Bottle or | 1 | | | Pre-Pierced Reseal Nebulizer* | A-Pack | 1 Each | | |
| tablets* | 4 UD Tabs A-Pack | 4 | | | | | | | |
| Atropine 1mg/10 mL | Syringe 10 mL | 3 | | | Blunt Cannula 18 g – 1 inch * | 18 G x 1 inch A-Pack | 5 2 | | |
| Calcium Chloride | Syringe 10 mL | 2 | | | Filter Needle | 18-21 G | 3 | | |
| 1 g/10 mL | , , | | | | Intranasal Mucosal | A-Pack | 1 Each | | |
| Ceftriaxone 2gm | 2gm vial | 1 | | | Atomization Device* | | 15.1 | | |
| Dextrose 50% 25 g/50 mL* | Syringe 50 mL A-Pack | 1 1 | | | Red Lock* | A-Pack | 1 Each | | |
| Diphenhydramine | Vial 1 mL | 2 | | | Replacement Form* | A-Pack | 1 | | |
| (Benadryl) 50 mg/1 mL Epinephrine | Amp/Vial 1 mL | 2 | | | Syringe 1mL (With needle/Luer Lock) | Syringe 1 mL | 5 | | |
| 1 mg/1 mL Epinephrine | Syringe 10 mL | 7 | | | Syringe 10 mL | Syringe 10 mL | 5 | | |
| 1 mg/10 mL | 25 1 1/1 1/10 | 2 | | | Syringe 20 mL | Syringe 20 mL | 1 | | |
| Ipratropium Bromide 0.02% (In Baggie)* | 2.5 mL Vial – UD A-Pack | 2 1 | | | | | _ | | |
| Ketorolac 15mg | 1ml Vial | 1 | | | Needle Pediatric Needle | 18 G x 1.5 inch 25 G x 1 inch | 2 | | |
| Lidocaine 100 mg/5 mL | Syringe 5 mL | 3 | | | 3 or 4-Way Stopcock* | 23 G X I IIICII | 1 Each | | |
| Magnesium Sulfate | Amp/Vial | 4 | | | Syringe w/ needle | Syringe 3 mL | 5 | | |
| 1 g/2 mL Methylprednisolone 125 mg | Vial | 1 | | | 3 mL-21/22 G x 1.5 inch* | A-Pack | 2 | | |
| Naloxone* | 4 x 2 mL Syringe | | | | | | | | |
| 2 mg/2 mL or | or | 4 | | | Replacing Hospital: | | | | |
| 0.4 mg/mL Drug Box | 2 x 10 mL Vial Total = 8 mg | 2 | | | | | | | |
| Nitroglycerin* | Bottle | 1 | | | MCA Medical Director's | s Name or post i | radio ordo | ering phy | sician: |
| 0.4 mg/tab | A-Pack | 1 | | | | • | | 01 / | |
| Ondansetron 2 mg/mL* | 2 mL vial | 2 | | | - | | | | |
| Ondansetron 4mg ODT* | 4mg tab | 2 | | | (Controlled Substance use of | only) PRINT NAM | 1E | | |
| Prednisone 50 mg tab* | 50 mg. tab A-Pack | 1 1 | | | Date: | | | | |
| Racepinephrine 2.25% 11.25 mg/0.5 mL | 0.5 mLVial & 3mL NS | 1 | | | | RAMEDIC'S STAT | | | |
| Sodium Bicarbonate 50 mEq/50 mL | Syringe 50 mL | 2 | | | SEM EMS Medication Bo above noted medication(s) |) used as prescrib | ed. I acce | ept pharm | acy seale |
| Sodium Chloride 0.9% | Vial 20-30 mL or | 1 | 1 | | SEM EMS Medication Bo | | seal | ed with b | reakaway |
| (Preservative free) | 10mL syringe | 2 | | | tag number | | | | |
| Sodium Chloride 0.9% | Bag 100 mL | 1 | | | Paramedic Signature: | | | Date: | |
| Tranexamic Acid (TXA) 100mg/ml | 10ml vial | 1 | | | RECEIVING PHARM The controlled substance (| | | | |
| | | OTM/ | DOGE | DOCE | Box number ha | s been reviewed. | The Supp | oly Use/Ro | eplaceme |
| CONTROLLED SUBSTANCES | UNIT/SIZE | QTY/ DOSE | DOSE GIVEN | DOSE WASTED | form reflects the C.S. cont | | | | |
| Fentanyl 50 mcg/ mL | Vial/Amp 2 mL | 3 | , | | administered by the Paran documented as administer | | | | |
| Midazolam 5 mg/1 mL | Vial 1 mL | 4 | | | documented as administer dosage form, volume, and | | | | |
| Morphine 10 mg/1 mL | Vial/Amp 1 mL | 2 | | | accuse form, volume, and | quantity per ivic | arear Com | | m, pone |
| Ketamine 100mg/ml | Vial 5ml | 1 | | | Name of Pharmacist on th | a Caali | | | |

Date: ______Hospital: _____

SEM A-PACK SUPPLY USE/REPLACEMENT FORM

| Date: Agency Name: | Unit | t #: Inc. # | : |
|--|-------------|---|-----------------------------------|
| Crew Names: | | | |
| Replacing Hospital: | | | |
| Paramedic's Statement | \neg [| MEDICATION | UNIT/SIZE |
| r aramedic 3 Statement | | Albuterol 2.5 mg/ 3 mL | Vial – UD 3 mL |
| SEM EMS A-Pack # has been opened and the noted medication(s) used as prescribed. I accept pharmacy seale | d | Aspirin 81 mg tablets | Chewable UD Tablets |
| SEM EMS A-Pack # sealed with breakaway # | | Dextrose 50% 25 g/50 mL | Syringe 50 mL |
| Patient Name:Patient DOB: | | Ipratropium Bromide 0.02% (In Baggie) | 2.5 mL Vial – UI |
| Paramedic Signature: Date: | _ | Naloxone 2 mg/2 mL or 0.4 mg/mL | 2 x 2 mL Syring 1 x 10 mL Vial |
| | $\dashv [$ | Nitroglycerin 0.4 mg/tab | Bottle |
| Replacing Pharmacist's Statement | | Ondansetron 2 mg/mL | 2 mL Vial |
| | | Ondansetron ODT | 4mg Tablet |
| The medication(s) in the sealed SEM EMS A-Pack | | Prednisone | 50 mg Tablet |
| #has been | | Nebulizer | |
| distributed according to the Medication/Use and Replacement | en t | Blunt Cannula | 18 G x1 inch |
| Policy of the participating MCA. All Medications are in the correct | | Intranasal Mucosal Atomization Device | |
| concentration, dosage, form, volume, amount, and not expired. | | Syringe w/needle 3 mL x 21/22 G x 1.5 inch | Syringe 3 mL |
| Signature of Replacing Pharmacist: | _] | 3 or 4-Way Stopcock Red Lock | |

| Paramedic's Statement | MEDICATION | UNIT/SIZE | QNTY | USED |
|---|---|---------------------------------------|------|------|
| Farametric 9 Statement | Albuterol 2.5 mg/ 3 mL | Vial – UD 3 mL | 6 | |
| SEM EMS A-Pack # has been opened and the noted medication(s) used as prescribed. I accept pharmacy sealed | Aspirin 81 mg tablets | Chewable UD Tablets | 4 | |
| SEM EMS A-Pack # sealed with breakaway # | Dextrose 50% 25 g/50 mL | Syringe 50 mL | 1 | |
| Patient Name:Patient DOB: | Ipratropium Bromide 0.02% (In Baggie) | 2.5 mL Vial – UD | 1 | |
| Paramedic Signature: Date: | Naloxone 2 mg/2 mL or 0.4 mg/mL | 2 x 2 mL Syringe or 1 x 10 mL Vial | 4 mg | |
| | Nitroglycerin 0.4 mg/tab | Bottle | 1 | |
| Replacing Pharmacist's Statement | Ondansetron 2 mg/mL | 2 mL Vial | 2 | |
| | Ondansetron ODT | 4mg Tablet | 2 | |
| The medication(s) in the sealed SEM EMS A-Pack | Prednisone | 50 mg Tablet | 1 | |
| #has been | Nebulizer | | 1 | |
| distributed according to the Medication/Use and Replacement | Blunt Cannula | 18 G x1 inch | 2 | |
| Policy of the participating MCA. All Medications are in the correct | Intranasal Mucosal Atomization Device | | 1 | |
| concentration, dosage, form, volume, amount, and not expired. | Syringe w/needle 3 mL x 21/22 G x 1.5 inch | Syringe 3 mL | 2 | |
| Signature of Replacing Pharmacist: | 3 or 4-Way Stopcock | | 1 | |
| Signature of Replacing Pharmacist. | Red Lock | | 1 | |
| Hospital: Date: | Replacement/Incident | Forms | 1ea | |